



climate change

Nowadays, when we talk about **climate change**, we are not referring to fluctuations in the Earth's overall climate as a result of natural factors, but rather to the hypothesis of **global warming** due to human causes. It is a matter of answering the question of whether the climate is changing—or has already changed—as a result of **human action**, and what effects that change will have.

The questions—the *frontiers of knowledge*—posed by this subject are very diverse. With each passing day, the emerging science of climate change more insistently asks (despite the ongoing existence of skeptics) what relation exists between the concentration of certain substances in the atmosphere and global warming. That warming can be understood as a result of the alteration of our **planet's energy equilibrium** by **greenhouse** gasses. Researchers are studying the functioning of this mechanism and its quantitative aspects, and are creating climate models that seek to calculate the effects of diverse scenarios involving the melting of the polar icecaps (and the glaciers), and rising sea levels. They study different alternatives for the geography of climate change (especially temperature and rainfall levels in different regions), and analyze the possible effects of those changes on the probability of catastrophes of climatic origin. They also explore the interaction of this science with the life sciences, studying the **biosphere's** vulnerability to global warming and its threat to **biodiversity**, analyzing its possible impact on different **ecosystems**, what species may be in most danger, how great that danger is, and how global warming may affect the availability of water and the capacity to grow food.

Scientists investigate the fundamental factors of this phenomenon, including the massive use of **fossil fuels** (major CO₂ emitters) as **energy sources**. But they also research the relation of global warming to other phenomena related to the intensive exploitation of the planet, including **deforestation**, land degradation, massive human occupation of the coasts and other sensitive areas, and so on.

In recent decades, awareness of climate change as a global problem requiring worldwide action has led to a proliferation of international organizations (of which the **IPCC** is one) and initiatives, which eventually led to the signing of the **Kyoto Protocol** (and more recently, the Bali Accord) for the reduction of greenhouse gasses. This strategy to **mitigate** climate change has had to

overcome the reticence of the most affected developed countries (which are the greatest emitters of such gasses) and consider how to distribute the load among developed and **emerging countries** (China is on its way to becoming the world's main producer of greenhouse gasses), dealing with its relation to development strategies and the **war on poverty**. In that sense, the main frontier lies in the design of concrete measures that must be adopted. Is the use of **alternative energy sources** the most promising way of fighting climate change? And of such sources, what role should the different renewable energies, such as solar and wind power, have? And what about nuclear energy? An especially pressing question today is whether the use of **bio-fuels** is affecting the cost of staple crops and thus helping to provoke an extremely serious food crisis?

Climate change again draws attention to the general problem of the relations between resources, population, and development—on a global scale, and for each geographic and social setting. In sum, it is a new facet of the problem of **sustainable development**. And all of this has important economic implications. From the viewpoint of developed countries, the **Stern Review** has approached this problem as a choice between alternatives whose costs can be compared. The author of that review defends a very clear option: the cost of climate change would be incomparably greater than the cost of fighting against it.

No matter what the costs and consequences of measures to mitigate it, we are also facing an ever-more-necessary preventative **adaptation**, not only to warming itself, but also to the effects of coastal occupation and hydrological imbalances. This adaptation is necessary to prepare us for the foreseeable multiplication of extreme climatic events (striking recent examples include hurricane *Katrina* and the *tsunami* in the Indian Ocean, which had devastating consequences for both developed and developing countries).

Today, these questions are at the forefront of political and media concern (with the noise, fury, and lack of rigor that often results from such concentration). The awarding of the 2007 Nobel Peace Prize jointly to the IPCC and Al Gore—who has become an apostle in the fight against global warming—merely emphasizes the importance of the challenge (which may also be an opportunity for humanity to rectify past errors) of climate change.



climate change on the planet earth

SERGIO ALONSO

*To my wife and companion, Mercedes, in memoriam.
To our daughters Aurora, Patricia and our son Carlos.*

Introduction

The year 2008 marks the twentieth anniversary of the establishment of the Intergovernmental Panel on Climate Change (IPCC). Its creation grew out of an agreement between the World Meteorological Organization (a part of the United Nations) and the United Nations Programme. Its goal was to supply independent scientific information—in principle, to politicians—about questions concerning climate change. Almost ten years earlier, in the first World Climate Conference, attention was drawn to the increase in human activities, indicating that they might produce climatic alterations on a regional and even a planetary scale. Some years later, the role of CO₂ in climate variations was evaluated, along with other gasses capable of contributing to the so-called greenhouse effect. There was also a call for objective, balanced, and internationally coordinated scientific judgment that would shed light on the consequences of an increased concentration of greenhouse gasses in the Earth's atmosphere, and the socio-economic effects they might produce. This environmental concern, which was officially made public about thirty years ago, although it was actually older, led to the establishment of the IPCC in 1988. In 2007, the Norwegian Nobel Committee decided that the Nobel Peace Prize should be "shared, in two equal parts, between the Intergovernmental Panel on Climate Change (IPCC) and Albert Arnold (Al) Gore Jr. for their

efforts to build up and disseminate greater knowledge about man-made climate change, and to lay the foundations for the measures that are needed to counteract such change"¹

Some of the terms appearing in this introduction will be dealt with in more detail further on, but some should be clearly defined from the start. First, we should point out that the planet undergoes climate change on a continuous basis. We can be certain that, in the past, the climate was different than it is now, and that it will continue to change in the future. At first, terminology was somewhat confusing, with a coexistence of terms such as climate variation, climate variability, climate change, and climatic change. Finally (and unfortunately) two meanings remain in use. In scientific terms, climate change, means any change undergone by the planet's climate, regardless of its cause. This option is used, for example, by the IPCC. However, the United Nations Framework Convention on Climate Change, which arose from the so-called Rio Summit of 1992, and the Kyoto Protocol (established following the Convention), use the same terminology to refer to climate change attributed directly or indirectly to human activity, which is superimposed on natural variability. Therefore, when climate change is mentioned, care must be taken to make it clear which of the two meanings is intended. Notice, for example, that the Nobel Foundation's declaration specifies that it is referring to

¹ This paragraph is taken from the official announcement of the awarding of the prize by the Nobel Foundation.

climate change induced by humanity. Later on, we will see that this Climate Change of anthropic origin can be explained in terms of an intensification of the natural greenhouse effect. That intensification derives from a change in the composition of our atmosphere brought about by human activity.

The contents of this contribution include, in the next section, the reasons why the planet's climate changes, whether natural or anthropic. In section 3, we will review recent observations of changes experienced by the climate. The following section will contain arguments based on numerical simulations of climate, which attribute those changes to human activity. Section 5 offers a few indications about the use of computer models to simulate the Earth's climate. On the basis of the trustworthiness of such computer models, section 6 deals with the generation of climate scenarios for the future. Our conclusions are presented in section 7, followed by the bibliography employed.

Why does the climate change?

The climate is dynamic, changing and even unrepeatable. It is the consequence of the energy the Earth receives from the Sun, and of the exchanges of energy among different parts of what is called the Climate System, which we can understand as a synonym for the Planet Earth. Those parts or subsystems are:

- The atmosphere, the planet's gaseous envelope, where we perceive the climate.
- The hydrosphere, consisting of oceans, seas, lakes, and so on.
- The lithosphere, the solid emerging crust of the continents, where we live.
- The biosphere, made up of all living beings, including mankind, and
- the cryosphere, which consists of all the ice that covers parts of the oceans and continents.

From a broad viewpoint, the climate can be defined as the state of the Climatic System, including its statistical properties. That is precisely what relates this definition of climate with the most classic and restricted one, which consists of a statistical description of environmental variables (for example, temperature, wind, surface humidity, and precipitation), using mean values and measurements of dispersion over long time periods, far superior to the typical periods of atmospheric weather.

The subsystems of the Climatic System mentioned above have very different dynamics. While some experience appreciable and continuous change (the atmosphere, for example, with its succession of quite different weather conditions—sunny, cloudy, windy, rainy, and so on), others change quite slowly, some so slowly

that their variability merits little consideration over the course of a single human lifetime, or even several generations (that would be the case of the lithosphere, for example, except for the most superficial layer). When the energy we receive from the Sun reaches the Earth, it is distributed among all the subsystems and is exchanged among them, establishing relations according to the dynamics of each. The differences among these exchanges give rise to the great variety of climates in different regions of our planet, which we know so well, and which are a manifestation of the climate's spatial variability.

But climate is also characterized by variability over time. The Sun's energy does not arrive in equal amounts at all times, nor do the subsystems of the Climatic System always behave exactly the same. Therefore, we should not expect the energy flows that occur to invariably coincide over time. In certain intervals of time, their statistics can coincide more or less, but there is no reason to think that this must always be that case.

Next, we will analyze in some detail the origin of variability, that is, what causes changes in the Earth's climate. Some of these causes are natural, others are not—meaning that they have to do with human activity. The extant level of knowledge about the mechanisms we will see below is generally high, but we must not forget that, whenever there is a lack of knowledge (and there always is, of course) there will be a certain degree of ignorance, which leads to uncertainty in the interpretation of the observed phenomena.

First of all, we must begin by speaking of the Sun and its relation with the Earth. Its energy travels through space as radiation (called solar or short-wave radiation). It reaches the Earth, which intercepts it no matter what part of its orbit it is in or what time of the year. Not all the energy intercepted is used by the Climatic System. A fraction of it (called albedo) is returned to space through different processes of reflection mainly by clouds and the Earth's surface. Planetary albedo is around 30%. Finally, the radiation that is not absorbed by the atmosphere reaches the surface, which heats up and, in turn, emits its own radiation (called terrestrial or long-wave radiation). A large part of that radiation is absorbed by the atmosphere, which then re-emits it, either towards the surface or upwards, thus returning energy to space. For the entire planet, in average terms over time, there is an overall balance of energy, but not in the planet's different parts, nor at all times. It is these specific differences that affect the climate (see Kiehl and Trenberth 1997).

But how can the balance of energy be altered? According to what has been said, there could be three reasons:

a) Changes in the energy intercepted by the Earth. These may be due to changes in the Sun's emissions of radiation as a result of solar activity itself, or to changes in the position of the Earth in its orbit around the Sun.

b) Changes in the Earth's albedo. These, then, would be due to cloudiness (both degrees of cloud cover and types of clouds), changes in the reflective properties of the ground (types of ground and vegetation), and changes in the particulate matter suspended in the atmosphere. These particles are known as "aerosols."

c) Changes in the flow of long-wave energy from Earth to space. In this case, the changes would be due to a modification of the absorbent properties of the atmosphere as a result of changes in its composition.

Changes in solar activity have been recorded. The most popular may well be what is called Maunder's Minimum, which is thought to have occurred between 1350 and 1850, coinciding with the so-called Little Ice Age (Hoyt, Schatten, and Nesme-Ribes 1994; Eddy 1976). Since that time it is estimated that radiation may have increased between 0.04% and 0.08%, with an increase of 0.05% between 1750 and the present (Wang, Lean, and Sheeley 2005).

But the Earth does not occupy a fixed position in relation to the Sun; it has a very approximate elliptical orbit—with the Sun at its focus—whose eccentricity changes cyclically over a period of about 100,000 years. That means that the Earth is not the same distance from the Sun, year by year, at the same point in its orbit—which is also changing. Moreover, the inclination of the Earth's axis with respect to the plane of its orbit (obliquity) is not constant. It is as if the Earth were a huge top, so the prolongation of its axis of rotation points to different places in the celestial dome in cycles lasting around 41,000 years. Also, the orbital ellipse changes its orientation in space, leading to what are called the precession of equinoxes. That means that the astronomical seasons take place in different parts of the orbit with cycles lasting approximately 19,000 and 23,000 years. The final result is that, even if the energy emitted by the Sun were constant, what actually affects the system varies, and is also distributed differently over the planet's surface. All of this constitutes what is called Milankovitch's Theory of Cycles, which, along with certain internal mechanisms, makes it possible to explain the succession of geological eras (Berger 1988).

The processes we have described are external to the Climatic System and in no way depend on human activity. Another possible cause of planetary Climate Change, which is also both external and natural but has no relation to the solar radiation received by the Earth, is the impact of meteorites or comets. This is something difficult to predict, but its consequences are important

when the objects are big enough. Their impact against the surface of the planet can cause a cloud of dust or water of such magnitude that incident solar radiation cannot reach the Earth's surface with the intensity it had before impact. In those conditions, the temperature can drop appreciably, leading to climate change. The extinction of some species, including dinosaurs, in what is called the K/T Boundary, seems to have this origin (Álvarez et al. 1981).

This cause, which we can qualify as exceptional, allows us to bring in those related with albedo. Following impact, there must have been a considerable increase in albedo because of the increased amount of aerosols (particulate matter) in the atmosphere. This would have reflected a very high fraction of solar radiation back into space. In consequence, the Climatic System would suddenly have had much less energy to heat the ground and, thus, the previous balance of radiation would have been altered. The result must have been a lowering of the temperature at ground level. Without reaching those extremes, something similar happens each time there is a volcanic eruption. Their effect on temperature has been observed following large eruptions and depends on the intensity of the eruption, and on how high up in the atmosphere the generated particles reach. The effect, which can last several years, has been widely studied (see, for example, Yang and Schlesinger 2002).

The aerosols we have considered up to now are of natural origin but, besides these, the Earth's atmosphere also contains many others stemming from human activity. Generally, they reduce air quality and many of them also lead to health problems. From a climatic standpoint they have two effects. One directly affects albedo, leading to lower temperatures. The other has an indirect effect, modifying the conditions in which clouds are formed and how long they last. The final result of this indirect effect is not well known. Nowadays, it is the subject of uncertainty.

Clouds' role in albedo depends on cloud cover, the type of cloud, and how long it lasts. Thus, high clouds (cirrostratus clouds, for example) allow solar radiation through, but absorb terrestrial radiation, while medium clouds (altocumulus clouds, for example) almost completely impeded the passage of solar radiation. The first case will result in a rise in temperatures, while in the second they will fall.

Albedo also depends, as mentioned above, on the reflective properties of the planet's surface. A frozen surface (high albedo, of 70% to 90%) is not the same as bare earth, prairie, or the ocean's surface (low albedo, <10%). Different types of terrain and ground-use mean that the climatic treatment of the Earth's surface is a complex problem and a source of uncertainty.

At this point, we cannot avoid commenting on one type of behavior that is characteristic of the Climatic System. Often, the effects of a process act on its own causes, generating a sort of cyclical, unending behavior called feedback. Feedback is typical in what are called non-linear or dynamic systems, and the Climatic System is one of them. The following example is relatively straightforward: let us suppose that, for whatever reason, the planet's surface temperature rises. One of the consequences will be the partial melting of its ice. Surface albedo will diminish, leading to decreased reflection of solar radiation. There will thus be more energy available to the system, and the temperature will rise further. The additional heating will lead to greater ice melting, reducing albedo even more, and so on and so forth. This, then, is a positive feedback cycle known as ice-albedo feedback. It was already identified in the nineteenth century (Croll 1890). In the Climatic System, there are many other positive feedback cycles like this one, but there are also negative ones. When those feedback processes act at the same time, it becomes very difficult to obtain detailed knowledge of the results, even though it is clear that they exist. The only possible way of dealing with the problem is through numerical simulation of those processes.

The last way of modifying the balance of radiation to be mentioned here might well have been the first: it is the main way of explaining the climate change the planet is experiencing today.

First, we will consider the role the atmosphere plays in exchanges of solar and terrestrial radiation, which is known as the Greenhouse Effect (GE). We have already mentioned that part of the radiation coming from the Sun—about 30%—is reflected back into space. If the Earth did not have an atmosphere, the planet's surface would have an average temperature of -18°C , barely enough to maintain the energy equilibrium between penetrating solar radiation and terrestrial radiation (infrared) that the Earth would emit at that temperature. The Moon, which has no atmosphere, has an average temperature like that. But since the Earth *does* have an atmosphere, things are radically different. The atmosphere's constituents absorb relatively little solar radiation (especially where there are no clouds) but some of them are very good at absorbing the infrared radiation emitted by the Earth and by the atmosphere itself. This leads to a warming of the lower layers of the atmosphere, which modifies the balance of radiation, reaching an average temperature of 15°C at ground level. This behavior by the atmosphere, which reacts differently to solar radiation than to terrestrial radiation, is the GE, whose name comes from its relative similarity to the behavior of such structures. The main cause of GE is water vapor (approximately 80% of the

total effect) and the second cause, at a considerable distance, is carbon dioxide (CO_2). The GE (to which the adjective "natural" is often added) is decisive in the planet's climate, which has allowed the existence of life, at least as we know it. The gasses that contribute to the GE are known as greenhouse gasses (GHG). That said, it should be obvious that the GE is also affected by aerosols and that the role of clouds can also be discussed in those terms.

Any change in the composition of the atmosphere, or in the concentration of its components, alters its properties of absorption, consequently altering the GE as well. The atmosphere's composition has been changing for as long as the Earth has existed. Nitrogen (N_2) and oxygen (O_2) predominate, although the major contributors to the GE are water vapor (whose concentration does not surpass 4% of the atmosphere's volume) and CO_2 (with a much smaller concentration, currently around 385ppm^2). If the atmosphere's composition changes, the GE will be modified and thus, the planet's mean surface temperature will change. Before the industrial revolution, the mean global concentration of carbon dioxide was around 280ppm , while it is now about 385ppm , as mentioned above. In these conditions, the planet's natural GE has been undergoing modification ever since the Industrial Revolution began. As the concentration of CO_2 has increased (those of other GHGs are also rising, including methane, nitrous oxide, CFCs, and so on) the GE has enhanced, more energy has become available in the lower layers of the atmosphere and, thus, conditions have arisen for warming on a planetary scale. This is not modern speculation; in the late nineteenth century, the Nobel scientist Svante Arrhenius estimated the effect of a 40% increase or decrease in atmospheric CO_2 on temperature, indicating that the glaciers could shrink or expand (Arrhenius 1896). Actually, by the end of the seventeenth century, there was already knowledge of the different behavior of certain substances with regard to solar and terrestrial radiation, which is the basis for GE.

By analyzing air from bubbles trapped in core samples extracted from polar ice, it is possible to obtain information about the evolution of the concentration of GHGs in past periods. These can also be compared to current levels. Figure 1 shows the value of carbon dioxide, nitrous oxide, and methane concentrations over the last 650,000 years. We can see that current values far surpass earlier ones, even in the warmer glacial periods. These are shown in Figure 1 as shaded bands. The lower part also shows variations in the concentration of deuterium, δD , in arctic ice. This serves as an indirect indicator of temperature variations. Note the values of δD in earlier warm




Figure 1. Variations of deuterium (δD) in antarctic ice, which is a proxy for local temperature, and the atmospheric concentrations of the greenhouse gases carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O) in air trapped within the ice cores and from recent atmospheric measurements. Data cover 650,000 years and the shaded bands indicate current and previous interglacial warm periods.

periods and in the present one, and the large difference in concentrations of GHGs. Unlike the present period, in which the relation between GHG and temperature is clearly established and the anthropic origin of the change in GHG is proven, there is still much to be discovered about many aspects of this relation in the past. It is thought that, in the Quaternary, changes in the concentration of CO_2 may have resulted from the simultaneous effects of biological and chemical processes in the ocean. They may also have been affected by changes in temperature (Köhler et al. 2005). Concentrations of CO_2 did sometimes surpass current levels in earlier periods, millions of years ago, but these are thought to have been the result of tectonic processes, such as volcanic activity, which determined changes of concentration (Ruddiman 1997).

Recently, as a result of the European EPICA research project, the time range has been increased to 800,000 years. The same conclusions hold with regard to concentrations of GHG indicated in the description of Figure 1 for the last 650,000 years (Lüthi et al. 2008; Loulergue et al. 2008).

Figure 2 shows variations in the concentration of CO_2 , CH_4 , and N_2O , but for shorter time periods (panels a, b, and c). The scale on the left of those panels shows the concentration of the corresponding GHGs, while the scale on the right represents what is called radiative forcing, which is equivalent to the intensification of the GE that implies increased concentrations of GHGs, as expressed in radiation units (Wm^{-2}). These three panels indicate that the change experienced by GHGs following the Industrial Revolution has no recent precedent: while the atmospheric concentration of CO_2 increased only 20ppm over the 8,000 years preceding industrialization, since 1750 it has increased more than 100ppm. Approximately two thirds of this increase is due to the burning of fossil fuels and the remaining third is due to land use change. Panel d represents the rate of change of the combined forcing of the same three GHGs, which gives an integrated value of 1.66Wm^{-2} since 1750. This amount is by far the greatest of all possible forcings associated with the different mechanisms responsible for climate change analyzed in this section.




Figure 2. The concentrations and radiative forcing by (a) carbon dioxide (CO_2), (b) methane (CH_4), (c) nitrous oxide (N_2O) and (d) the rate of change in their combined radiative forcing over the last 20,000 years reconstructed from antarctic and Greenland ice and firn data (symbols) and direct atmospheric measurements (panels a,b,c, red lines). The grey bars show the reconstructed ranges of natural variability for the past 650,000 years. The rate of change in radiative forcing (panel d, black line) has been computed from spline fits to the concentration data. The width of the age spread in the ice data varies from about 20 years for sites with a high accumulation of snow such as Law Dome, Antarctica, to about 200 years for low-accumulation sites such as Dome C, Antarctica. The arrow shows the peak in the rate of change in radiative forcing that would result if the anthropogenic signals of CO_2 , CH_4 , and N_2O had been smoothed corresponding to conditions at the low-accumulation Dome C site. The negative rate of change in forcing around 1600 shown in the higher-resolution inset in panel d results from a CO_2 decrease of about 10 ppm in the Law Dome record.

In essence, what we have presented so far are the climate drivers related to the balance of radiation on a global scale. As was indicated above, the climate is a consequence of energy flows in different parts of the Climatic System. Now is when a large number of processes with their own internal dynamics come into play, with a great wealth of time scales, making the system truly complex. As a result, the Climatic System is very difficult to deal with. To study it in its entirety calls for numerical simulation. What must be clear is that whenever the functioning of one part of the machinery is modified, the end result will be a change of climate (see IPCC 2007).

Nowadays, when we speak of (human-induced) climate change, we are referring to climate change

observed today, which is a consequence of the intensification of the GE. In the final analysis, this is a change in how Planet Earth functions as a consequence of human activity. That is what has come to be known as Global Change, leading some researchers, including Nobel Scientist Paul Crutzen, to say that the planet has entered a new era characterized by anthropic impact. That is why they propose this era be called the "Anthropocene" (Crutzen and Stoermer 2000).

Observing change

But is the climate really changing? Many people wonder, and ask the specialists. From a general viewpoint, the answer is yes. This planet's climate has always been changing. And now? In the Anthropocene?

Yes, now too. There are two aspects of current climate change that should be mentioned. The first is that, unlike earlier change, it has such a short time scale that change is appreciable over a period comparable to a human lifetime. The second is that humanity has never before had the capacity to interfere with climate on a global scale. It so happens that this planet's climate made life, including human life, possible. Now, the human species is capable of modifying that climate. These two characteristics make it possible to state that, strictly speaking, there is no past precedent for current climate change.

In this section, we will offer some of the evidence for current climate change. The following one will deal with procedures that have led to the conclusion that human activity is responsible for the observed changes.

In its fourth, and most recent report (IPCC 2007), the IPCC indicates that, compared to its third report (the third is designated by the acronym, TAR, and the fourth, AR4), there are now better data bases, more evidence, greater geographic coverage and a better understanding of uncertainties. As a result, AR4 indicates that the warming of the Climatic System is unequivocal, as can be deduced from observations of increased mean atmospheric and oceanic temperatures on a planetary scale, extensive melting of snow and ice and the global rise in the average sea level.

In TAR, calculations of warming of the mean global air temperature at ground level between 1901 and 2000 gave a linear trend of $0.6 \pm 0.2^\circ\text{C}$ per century. This was surpassed by AR4's calculations of that rate for the period between 1906 and 2005, which was $0.74 \pm 0.18^\circ\text{C}$ per century. The acceleration of warming becomes even clearer when we use only the last fifty of those one hundred years (1956–2005), and even more so in the last 25. In those cases, the resultant linear trend is $1.28 \pm 0.26^\circ\text{C}$ per century and $1.77 \pm 0.52^\circ\text{C}$ per century, respectively.³ The temperature increases noted here are very likely unprecedented on Earth, at least in the last 16,000 years.

Changes in temperature extremes have also been observed, and these are consistent with warming of the lower layers of the atmosphere. Thus, the number of cold and frosty nights has diminished, while the number of warm days and nights, and heat waves, has increased.

If we analyze the spatial distribution of these trends (which are greater on land than over the oceans) and the seasonal values, we will find important differences. The same occurs with separate calculations of maximum and minimum temperature trends. For example, the results of an analysis of temperature trends on the Balearic Islands over a thirty-year period ending in 2006 (OCLIB 2007) showed a linear trend of $4.83 \pm 1.85^\circ\text{C}$ per century for the

maximum temperature, with $5.14 \pm 1.89^\circ\text{C}$ per century for the minimum. The maximum value for the minimum temperature appeared in the summer ($8.01 \pm 3.17^\circ\text{C}$ per century), while the maximum value for the maximum temperature ($7.99 \pm 3.01^\circ\text{C}$ per century) appeared in the spring. It is important to note the large differences encountered here with respect to global values, even with the highest one quoted before, which corresponds to a period of 25 years.

The average temperature of the ocean has also risen, at least to depths of about 3,000 meters. It is estimated that, since 1955, the ocean has absorbed around 80% of the excess heat resulting from the GE. This results in the expansion of seawater and significantly contributes to sea level rise.⁴

Moreover, we must point out important changes in the cryosphere. For example, the surface area of arctic sea ice has diminished an average of 2.7% per decade, and that reduction process intensifies in northern hemisphere summers, where it reaches 7.4%. In the summer of 2007, the reduction of surfaces with at least 15% ice coverage was especially notable, after AR4 were developed. Such covered surfaces reached a summer minimum of 7.5 million square kilometers (averaged between 1979 and 2000) while, in the summer of 2007, only 4 million square kilometers were covered. That is the smallest surface area since Earth-observation satellites have existed. Values for the summer of 2008 show a slight recovery compared to 2007, but still far below the previously indicated average.⁵

Figure 3 indicates observed changes in the last century-and-a-half in the mean global surface temperature (panel a), the average sea level (panel b) and the surface of the Northern hemisphere covered with snow (panel c). The relative scale at the left of Figure 3 shows the variation of those changes with respect to the average value between 1961 and 1990.

Global rainfall measurements are also being affected by current climatic change. To start with it must be said that there has been a continuous increase in the total content of water vapor in the atmosphere, which is coherent with the temperature increase in the troposphere. Precipitation has been modified to an unequal extent in different geographic areas. While it has significantly increased in eastern parts of North and South America, northern Europe, and northern and central Asia, the climate is now drier in the Sahel, the Mediterranean, Southern Africa, and part of Southern Asia. If we look at the extremes, on one hand the occurrence of strong rains over land has become more frequent, but on the other more intense and lasting droughts have been observed since the nineteen seventies, particularly in the tropics and subtropics,

³ Warming has been observed in the average global temperature at surface level and in the troposphere. At higher levels—the stratosphere, for example—cooling of between 0.3°C and 0.9°C per decade has been observed since 1979, although this has diminished in recent years.

⁴ Variation in sea level is a complex problem lying outside the scope of this text. From a climatic standpoint, the main contributions, in almost equal measure, are the expansion of sea water (including the salinity effect) and the melting of continental ice. On geological time scales, there have been very important changes in sea levels. For example, it is estimated that, during the ice ages, level was over 100 meters lower than today.

⁵ Information drawn from <http://nsidc.org/arcticseaicenews/index.html>, consulted on 17 August, 2008.

sometimes in combination with flooding in those same geographical areas.

It is difficult to obtain figures on the trends of global precipitation, due mostly to the characteristic discontinuity of the variable and to measurement methods. As an example on a much smaller scale, the results offered below are from an analysis of precipitation trends in the Balearic Islands over a series of 55 years through 2006 (OCLIB 2007). Smoothing the annual precipitation series with a 5-year filter generates a tendency of $-170 \pm 123\text{mm}$ per century, which becomes $-192 \pm 38\text{mm}$ per century when the annual series is filtered with a 30-year average. Consideration must be given to the fact that the normal precipitation in the Balearic Islands is close to 600mm per year, which represents a decrease in rainfall tending towards 30% in




Figure 3. Observed changes in (a) global average surface temperature; (b) global average sea level from tide gauge (blue) and satellite (red) data and (c) Northern Hemisphere snow cover for March–April. All differences are relative to corresponding averages for the period 1961–1990. Smoothed curves represent decadal averaged values while circles show yearly values. The shaded areas are the uncertainty intervals estimated from a comprehensive analysis of known uncertainties (a and b) and from the time series (c).

one hundred years. This reduction has not been equally spread among the seasons, nor for all types of precipitation. Decreases have been greater in fall and winter, and much less so in spring and summer, linked to a decrease in the number of days with moderate rainfall, although the number of days with weak rainfall has increased, as has the number of days with strong rains, though to a lesser degree.

Observed changes in rainfall data are explained, in part, by the previously mentioned increase in atmospheric water vapor content, but also by the change in patterns of atmospheric circulation characteristic of the climate's natural variability. These include North Atlantic Oscillation (NAO) and the phenomenon El Niño/Southern Oscillation (ENSO).

Scientists are also confident about changes observed in some other extreme phenomena not mentioned here (for example, increases in the number and intensity of tropical Atlantic cyclones). But for others (tornados, lightening, hail, Antarctic sea ice, and dust storms) there are not yet enough reliable results to allow us to be certain that they have experienced variation in the present climate.

For more information on the changes observed it is necessary to consult AR4 (IPCC 2007).

Attribution of observed climate change

The term "attribution" is used here to indicate the process by which we evaluate whether the observed changes are consistent with quantitative answers to the different causes of planetary climate change simulated with well-tested models, and not consistent with other physically possible alternative explanations. In this section, we will take it for granted that the climate can be simulated in a sufficiently adequate manner; in the following one, we will try to offer arguments that make it clear that this is indeed the case.

Ever since the IPCC drew up its first report in 1990, the subject of attribution has been addressed. In that first report (FAR) there was not sufficient observational data on anthropic effects on climate. The second report (SAR) concluded that overall evidence suggested a discernible human influence on the twentieth century's climate. TAR indicated that the greater part of warming observed in the last 50 years was probably due to increased concentration of GHGs. Since that report, confidence in the evaluation of humanity's effect on climate change has increased considerably. We have more evidence, and the methodology of attribution has improved. All of this appears in AR4 and will be summarized below.

Attribution of current climate change will be carried out here using the results for temperature, the variable most clearly determined, and whose simulation is most

resolved. The observed evolution of temperature will be compared with what models are able to simulate. Figure 4 offers the results of a comparison of mean global temperatures with what climate models simulate for the twentieth century in different circumstances. In both panel a and panel b, the black curve represents the evolution of the global mean surface temperature. The values deduced from the scale on the left are temperature differences with respect to the average in the period 1901–1950. The red curve on panel a represents the mean evolution of the simulated temperature. It is obtained by averaging out the results of each of the individual models, whose different processes are represented in other. For this simulation, the models include known causes of climate change—specifically, natural ones, including volcanic eruptions—and those that are a consequence of human activity, using the known evolution of atmospheric concentrations of GHGs and aerosols. The result of this attribution experiment can be summed up by saying that there is a strong correlation between the evolution of observed and simulated temperatures, that the envelope of individual simulations almost completely includes the curve of observations, and that the average of the models closely approximates that of the observations when conveniently filtered by a time average (not shown in the figure).

Panel b presents the results of simulating the evolution of temperature using only natural causes of climate change. As before, it shows both the individual processes of models, in light blue, and the average of all the simulations, in darker blue. But here, the same conclusions cannot be drawn. The natural forcings can only explain the evolution of temperature through approximately the middle of the past century. In fact, a comparison of the two panels does not reveal large differences in the two simulations for that time period. The differences arise in the second half of the twentieth century. It is necessary to include anthropic causes in the simulations in order to explain the temperature trend in the second half.

This type of experiment had already been carried out in TAR (IPCC 2001), but the conclusions were not as trustworthy as in AR4. Moreover, equivalent studies have now been carried out for the different continents, separately for land and sea, and for other different temperature variables. The results are coherent with what has been stated above.

Climate research should always tend to reduce uncertainty while also achieving increasing realism in its simulations. Figure 4 shows important discrepancies between the simulations and mean surface temperature calculated by direct measurements around 1940. Analysis

of the origin of the temperature observations concludes that there is a bias in the observed values as a result of the method employed to measure the sea's surface temperature, which obviously forms a part of the planet's surface temperature (Thompson et al. 2008). If the observed values were corrected, the discrepancy would be reduced, bringing the observed temperature evolution closer to the simulation. At the time that AR4 was published, the above was not yet known, but the results were still considered sufficiently realistic to indicate that "most of the observed increase in global average temperatures since the mid-20th century is very likely⁶ due to the observed increase in anthropogenic GHG concentrations."

Simulation of the Earth's climate with models

Knowledge of the mechanisms that determine climate, set out in section 2, is partial but sufficient to allow us to simulate it (not in a laboratory, of course, but using complex models run by powerful computers). It has become possible to reproduce the current and past climates with sufficient accuracy, as well as the fundamental known traits of the climate in far earlier geological eras. Thanks to that, attribution exercises have been carried out, as indicated in section 4, and we can also think about inferring the possibilities of the future climate, including man's role in it. This last matter will be addressed in the following section.

Let us now look at climate simulation models in some detail. In the first place, we should state that such models are not an invention of climate researchers; in physics and other sciences, models are generally employed and they have turned out to be extraordinarily useful for advancing knowledge. In general terms, a model is a simplification of reality used as a tool to describe and explain phenomena of scientific interest. Models are sometimes constructed through mathematical equations that describe empirical relations among variables characteristic of the system being studied. For example, such relations can be obtained on the basis of an adequate statistical treatment of those variables. At other times, previous and independently established physical laws are used to establish the relations among variables. Moreover, in this case, they allow an interpretation of why this relation exists because, in fact, that is what these laws express. Finally, there are also mathematical equations that relate variables but are in this case based on physical laws.

In all cases, a set of equations is obtained that makes it possible to offer an approximate (remember, these models are simplifications) description of reality. It is precisely this fact that makes it possible to at least partially explain the discrepancies that appear between

⁶ IPCC uses this term to indicate that the probability surpasses 90%.



a simulated description of reality generated by a model, and the reality of observations of a real phenomenon.

Once the set of equations that constitute a model is obtained, those equations must be written in such a way as to furnish quantitative information about the system being studied. In the case we are discussing here, at the very least, they would have to furnish values for temperature and precipitation in order to reveal the fundamental traits of climate. Moreover, they would have to do so for the entire planet and, actually, at different levels of the atmosphere,




Figure 4. (a) Global mean surface temperature anomalies relative to the period 1901 to 1950, as observed (black line) and as obtained from simulations with both anthropogenic and natural forcings. The thick red curve shows the multi-model ensemble mean and the thin lighter red curves show the individual simulations. Vertical grey lines indicate the timing of major volcanic events. (b) As in (a), except that the simulated global mean temperature anomalies are for natural forcings only. The thick blue curve shows the multi-model ensemble mean and the thin lighter blue curves show individual simulations. Each simulation was sampled so that coverage corresponds to that of the observations.

from the lowest ones, at ground or sea level, to the highest. And that is only the part that deals with the atmosphere, because in other subsystems, it will be necessary to know many other variables (for example, the salinity and temperature of the oceans, ice mass, and the properties of soil and vegetation), at different levels or depths as well. The conclusion we must draw from all this is that the model's equations must be applied to a large number of points in space. Many mathematical operations have to be carried out in order to determine all the variables that describe the state of the Climatic System at a single instant in time. But in order to characterize climate, we must know what happens, not only at a specific moment, but over the course of sufficiently long time spans. That is, an enormous succession of individual instants.

How can we approach such a huge task? The answer is not immediate. First of all, if we want to obtain useful climate information in a reasonable time, we must use very powerful computers—the most powerful in the world. In order to do so, we must, again, simply the model, writing it in a form that is adequate for computer work. Once this is done, computers will be used to carry out the millions and millions of mathematical operations needed to obtain climate simulations for various decades, centuries, and so on, in a reasonable amount of time. Numerical simulations of climate are often mentioned in order to designate the means by which the desired climatic information is obtained.

The most advanced models of climatic simulation include formulas that address processes in the atmosphere, the oceans, the Earth's surface, and the cryosphere, atmospheric chemistry and the modeling of aerosols. They also deal in a linked way with atmosphere-oceanic interactions. Some models include mechanisms for maintaining energy flows at reasonable values, but nowadays, due to advances in research, most of them do not need this adjustment because the flows obtained directly by the simulations are already realistic. Those climate simulation models that include equations for the treatment of the processes mentioned here are generically called Atmosphere/Ocean General Circulation Models (AOGCMs). Many models exist, generally linked to leading research centers around the world, and their climate simulations offer different results, all of which are plausible. There are intercomparison projects and programmes in which results are contrasted in order to verify performance, which also makes it possible to establish confidence levels for the results. The IPCC itself bases a large part of its evaluation reports (see chapters 8 and 9 of AR4, IPCC 2007) on simulations. Confidence in climate simulation has been obtained by verifying that

the results are sufficiently realistic in comparison with observations. Those results affect the different subsystems of the Climatic System and known of variability modes of the current climate, including the phenomena of El Niño/Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO), patterns of anticyclonic blockage and the variability of monsoons. But verification by contrast with the present climate is not the only source of confidence. From a conceptual viewpoint, the primary source is that these models utilize physical laws that were independently established before the problem of climate simulation was even addressed. Moreover, it has become possible to simulate important traits of the climate of the last 2,000 years, as well as earlier climate change, such as the warm period in the Holocene, some 6,000 years ago, and the variability of the ice ages. It goes without saying that the results are reliable enough to foster confidence in the use of such models, despite the fact that there are still areas of uncertainty.

One of the main advantages to using models to simulate climate is that processes included in those models can be activated or deactivated at will. It is enough to eliminate the set of equations that affect a specific process in a given model. That model is then capable of simulating the planet's climate with or without the activity of the process (or processes) under study. Thus, for example, following a volcanic eruption,




Figure 5. Left Panel: Global GHG emissions (in GtCO₂-eq) in the absence of climate policies: six illustrative SRES marker scenarios (coloured lines) and the 80th percentile range of recent scenarios published since SRES (post-SRES) (gray shaded area). Dashed lines show the full range of post-SRES scenarios. The emissions include CO₂, CH₄, N₂O and F-gases. Right Panel: Solid lines are multi-model global averages of surface warming for scenarios A2, A1B and B1, shown as continuations of the 20th-century simulations. These projections also take into account emissions of short-lived GHGs and aerosols. The pink line is not a scenario, but is for Atmosphere-Ocean General Circulation Model (AOGCM) simulations where atmospheric concentrations are held constant at year 2000 values. The bars at the right of the figure indicate the best estimate (solid line within each bar) and the likely range assessed for the six SRES marker scenarios at 2090-2099. All temperatures are relative to the period 1980-1999.

the additional effect of expelled aerosols can be included, or the intensification of the GE can be eliminated while pre-industrial concentrations of GHGs are being considered. That, precisely, is the basis for the attribution of climate change dealt with in the previous section.

If we do not want to use large computers, or do not have access to them, there are also more modest solutions, which are not necessarily less useful. It is possible to gain access to a second level of climate simulation using a new simplification of the Climatic System. In other words, it is possible to simplify the complexity of the model—which is already, itself, a simplification of reality—in order to be able to work with personal computers or the like. In such cases, it is a matter of making sure that the simple models offer simulations that are compatible with those being carried out with AOGCMs.

To give us an idea: at the maximum extreme of simplification, we could consider the Earth a sphere that receives energy from the Sun and maintains the equilibrium of that energy with the energy it reflects, and that which the Earth itself radiates into space. In such conditions, a temperature—called the equilibrium temperature—is determined. It turns out to be around -18°C and is very different than the mean temperature on Earth, which is about 15°C. These same figures were mentioned above when discussing the natural GE. In other words, the equilibrium temperature is obtained by a maximum simplification of the system (specifically, by eliminating the atmosphere), which makes conditions more similar to those on the Moon than on the Earth. Including the atmosphere allows us to assign a temperature increase of some 33°C to the GE. If we really think about it, we realize that is a spectacular amount, especially when compared to what is thought to be the temperature oscillation associated with geological eras or abrupt climate changes. None of these is even half the amount indicated for warming due to natural GE (Masson-Delmotte et al. 2005).

With other simple models—though less simple than the one described above—it is possible to calculate the distribution of the equilibrium temperature for different latitudes on Earth, to establish elemental considerations about the clouds' role, and to determine other potential climates when all the ice has melted, or when the Earth is totally covered with ice, as well as the transitions between such states, and so on. One advantage of simple models, compared with more complex ones, is that they allow us to carry out a large number of different experiments—because they need much less time to resolve the equations than more complex models.

Climate projection for the future

It is important to emphasize that climate models are the most important, if not the only, tools for carrying out simulations of the planet's climate. In order to be able to use them with any sort of guarantee, experiments have been carried out to reproduce the present climate and the past climate, and to explain the climate change being experienced by the Earth. Since the basic equations come from physical laws and the simulation is realistic, there is great confidence in the use of such models. Clearly, there are still aspects to be discovered with regard to how the Climatic System functions, and this lack of knowledge produces uncertainty. Nevertheless, by accepting the results of the simulation when they are verified by observation, we are indicating that the knowledge we already possess about how that System works is sufficient, and what is still unknown would not be able to substantially modify the simulations. If that were not the case, that is, if our ignorance implied important consequences for such simulations, research would already have detected it.

7

GtC are gigatons of carbon, that is, a thousand million tons of carbon.




Figure 6. Projected surface temperature changes for the early and late 21st century relative to the period 1980 to 1999. The left and right panels show the AOGCM multi-model average projections ($^{\circ}\text{C}$) for the B1 (top), A1B (middle) and A2 (bottom) SRES scenarios averaged over the decades 2020 to 2029 (left) and 2090 to 2099 (right).

That being said, it should be clear that simulation of the present climate is not the same problem as simulation of the future climate. In the first case, we know what changes took place in the past, leading up to the present. We know how radiation intercepted by the Earth has changed, and we know how the atmospheric composition has changed—not only with regard to the concentration of GHGs but also, for example, to volcanic eruptions. The forcing of models with real and known conditions has made it possible to reconstruct the present climate. But from now on, we do not know what the conditions of the Earth's atmosphere will be, yet that knowledge is imperative if we are to simulate the future climate.

We know from the past, for example, that annual emissions of CO_2 of fossil origin have increased from an average of 6.4GtC^7 per year in the nineteen nineties, to 7.2GtC per year between 2000 and 2005. These emissions, along with those of the past, have partially determined the concentration of CO_2 in the atmosphere, just as other processes have done with other GHGs. The problem of determining the concentration of GHGs on the basis of emissions is not a simple one: it is necessary, once again, to resort to simulation using models. In this case, they are models of the cycles of carbon and other elements. It is necessary, for example, to take into account how carbon is fixed in the soil and in the seas ("carbon sinks"), which in turn depends on many factors.

Supposing that this problem is resolved, it will still be necessary to know how future GHG emissions evolve. What will definitely be clear by now is that this depends on many conditions, most of which are fundamentally socioeconomic in character and difficult to determine. In response, work is being done with different plausible hypotheses generally called scenarios. Ever since the earliest IPCC reports (FAR and SAR), attention has been paid to defining emissions scenarios, which were initially included in the reports themselves. Following the second report, however, specific work on scenarios was commissioned (IPCC 2000), which generated those scenarios currently being used to project the climate into the future. They are called SRES, an acronym that reflects the character and title of the work: Special Report on Emissions Scenarios.

In a nutshell, work is being done with four storylines (A1, A2, B1, and B2) conditioned by "forces" such as population, economy, technology, energy, agriculture, and soil use. In A1 and A2 more weight is given to economic growth, while in B1 and B2 environmental aspects take the fore. Also, whereas A1 and B1 project on the basis of a globalized world, A2 and B2 emphasize regional and local solutions. Each of these lines generates different scenarios, making a total of




Figure 7. Relative changes in precipitation (in percent) for the period 2090–2099, relative to 1980–1999. Values are multi-model averages based on the SRES A1B scenario for December to February (left) and June to August (right). White areas are where less than 66% of the models agree in the sign of the change and stippled areas are where more than 90% of the models agree in the sign of the change.

40. Normally, these are organized as families, coinciding with the name of their lines, except for A1, which has the following breakdown:

- A1FI, with intensive use of fossil fuels,
- A1T, with use of non-fossil energy sources,
- A1B, with a balanced use of different sources.

Clearly, we do not know what path humanity will take from here on, so all of those scenarios are considered equally probable.

Each of these SRES emissions scenarios is associated with concrete GHG emissions values over the course of the twenty-first century. Then, using adequate models, future concentrations of GHG are deduced, and the future evolution of those concentrations allows us to project the climate into the future, thanks to climate simulation models. The result is a group of climate projections for each of the SRES being considered. Because they differ from the climatic conditions set as reference, they lead to different future scenarios of climate change. Those scenarios or projections can be global, or limited to specific regions of the world's geography.

The left panel of Figure 5 shows the evolution of GHG emissions during the twenty-first century. The figure includes emissions of all GHG in what is called equivalent CO_2 . In calculating this, account is taken of the same GE intensification effect as all the GHGs being considered. As well as all the SRES scenarios described above, results are given here for other scenarios that appeared after the Special Report (IPCC 2000) was published. These modify the contribution of

certain "forces" that affect the storylines being considered. The right panel shows projected mean surface temperatures for various families of scenarios and the projection that would correspond to no increase of GHG over the amounts registered in the year 2000. It should be pointed out that, even if that were the case, the temperature would continue to rise, though at a much slower rate.

An analysis of projections for the first two decades of this century offers results that depend very little on the scenario being considered and the model employed (0.2°C per decade). That is not the case, however, for the final decades of the century; they strongly depend on the scenario being considered, and also on the model employed. For example, the mean multi-model estimation for scenario B1 at the end of the century is 1.8°C (probably with a range of 1.1°C to 2.9°C), while for scenario A1FI, it is 4.0°C (probably with a range of 2.4°C to 6.4°C), which is always higher than the mean for the period from 1980 to 1999. Note that those values are far above those observed for the increase in mean surface temperature during the twentieth century.

These temperature projections have been used to evaluate the effect on the global average sea level (including the contribution of ice melting in Greenland and Antarctica as well). The rise at the end of the twenty-first century—depending on which scenario is chosen, of course—would lie between a minimum of 0.18 to 0.38 meters for scenario B1 and a maximum of 0.26 to 0.59 meters for scenario A1FI. Those values are relative to the global average sea level between 1980 and 1999.

The AOGCM models make it possible to carry out global climate projections in which spatial and temporal variability is apparent. AR4 includes many such projections (see IPCC 2007, chapter 10), only a few of which are presented here. Figure 6 shows maps of multi-model mean surface temperatures with a clear predominance of values in the arctic region, where the temperature could increase by more than 7°C by the end of the century. In general, the projected warming for the twenty-first century is expected to be greater over land and at higher latitudes of the northern hemisphere, and lesser over the South Seas and part of the North Atlantic.

Figure 7 shows projections for seasonal rainfall. While global amounts are expected to rise, in the majority of terrestrial sub-tropical regions they will probably decrease. In the upper latitudes, precipitation will probably be greater.

Projections for other important aspects of the climate have also been obtained. Generally, it could be said that all of them continue the tendencies observed in the twentieth century but most show increases in those tendencies.

Special mention should be made of the melting of ice in Greenland, although the time scale is more than a century. Some 125,000 years ago, the temperature in the North Atlantic zone remained higher than at present for a prolonged period of time. The reduction of the ice mass led the sea levels to rise between 4 and 6 meters. Now, if the temperature remained between 1.9 and 4.6°C higher than pre-industrial levels for at least a thousand years, melting Greenland ice would cause a rise in planetary sea levels of 7 m.

One of the most important applications of climate projections is the analysis of the consequences of climate change or, as it is generally called, the impact of climate change. This has considerable social importance because its effects are local. In order to determine them, it is necessary to have climatic projections with much greater resolution than those offered by global models. This is done with different methodologies and is generally called "downscaling." One of the most common employs regional-scale models nested in global models and run in a linked, simultaneous way. That is dynamic downscaling. Another possibility involves using empirical statistical relations that have been determined for the present climate—they are supposed to remain valid in the future—in order to gain resolution on the basis of future climate projections obtained with AOGCM. There are also methodologies that combine the two mentioned above. More information on downscaling is available in chapter 11 of AR4 (IPCC 2007).

Conclusions

During the Anthropocene, planet Earth is experiencing a change of climate that, strictly speaking, has no precedent in the past. The burning of fossil fuels and general human activity have modified the atmosphere's composition, increasing the concentration of GHGs to levels never before attained, at least in the last 800,000 years. The GE, which has allowed life to exist on Earth, is being intensified anthropogenically, leading to an increase in the global mean surface temperature in the twentieth century that has no antecedents, at least in the last 16,000 years. Along with this change of temperature, a rise in sea levels has also been observed, as well as the reduction of snow coverage on the continents, and sea ice in the Arctic Ocean. Moreover, climate patterns are changing, including rainfall, NAO, and the phenomenon ENSO, among others. The frequency with which certain extreme phenomena occur is also changing.

If GHG emissions continue at the current rate, observed climate change will accelerate in the present century. In fact, even if the concentrations of those gases remained at their current levels, temperature increases and the resulting effects would continue to occur for decades, though with lesser intensity.

The social and economic consequences of the changes observed have already become significant in some zones (changes of habitat, exhaustion of certain species' capacity to adapt, modification of crop seasons, problems with water resources, changes in the distribution and occurrence of certain diseases, and so on), but it is believed that they will become even more significant as warming intensifies. From a human standpoint, the most disadvantaged societies, with the lowest levels of development, will be the most vulnerable.

Global warming can no longer be stopped; we are already suffering the consequences of what we began with the Industrial Revolution. It is clear that we have to reduce emissions, and that is intrinsically good for the environment in general. But we must also strive to adapt to the coming climate and understand that, beyond living with a certain level of risk, it will be necessary to face the cost of adapting. At any rate, that will be much less than the cost of doing nothing. Policymakers have to play their role and the society also the own. Obviously, as members of society, scientists, too, must participate. Research must be intensified, eliminating doubts, improving climate projections, offering clues as to how to reduce climatic vulnerability and risk, and seeking out more efficient means of energy use, less contaminating systems, and so on.

We will undoubtedly have to make some slight changes of lifestyle so that developing countries can attain an adequate level of wellbeing. The future humanity expects nothing less of us.

Bibliography

- Álvarez, L. W., W. Álvarez, F. Asaro, and H. V. Michel. "Asteroid Extinction Hypothesis." *Science* 211, 1981, 654–656.
- Arrhenius, S. "On the influence of carbonic acid in the air upon the temperature on the ground." *Philos. Mag.* 41, 1896, 237–276.
- Berger, A. "Milankovitch theory and climate." *Reviews of Geophysics* 26 (1988): 624–657.
- Croll, J. *Climate and Time in Their Geological Relations: A Theory of Secular Changes of the Earth's Climate*. 2nd ed. New York: Appleton, 1890.
- Crutzen, P., and E. F. Stoermer. "The 'Anthropocene.'" *Global Change Newsletter* 41, 2000, 12–13.
- Eddy, J. A. "The Maunder Minimum." *Science* 192, 1976, 1189–1202.
- Hoyt, D. V., K. H. Schatten, and E. NESMES-RIBES. "The hundredth year of Rudolf Wolf's death: Do we have the correct reconstruction of solar activity?" *Geophys. Res. Lett.* 21, 1994, 2067–2070.
- IPCC. *Special Report on Emissions Scenarios*. Nakicenovic, N., and R. Swart (eds.). Cambridge and New York: Cambridge University Press, 2000.
- IPCC. *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Houghton, J. T., Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Da, K. Maskell, and C. A. Johnson (eds.). Cambridge and New York: Cambridge University Press, 2001.
- IPCC. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller (eds.). Cambridge and New York: Cambridge University Press, 2007.
- Kiehl, J., and K. Trenberth. "Earth's annual global mean energy budget." *Bull. Am. Meteorol. Soc.* 78, 1997, 197–206.
- Köhler, P., F. Joos, S. Gerber, and R. Knutti. "Simulating changes in vegetation distribution, land carbon storage, and atmospheric CO₂ in response to a collapse of the North Atlantic thermohaline circulation." *Clim. Dyn.* 25, 2005, 689–708.
- Louergue, L., A. Schilt, R. Spahni, V. Masson-Delmotte, T. Blunier, B. Lemieux, J.-M. Barnola, D. Raynaud, T. F. Stocker and J. Chappellaz. "Orbital and millennial-scale features of atmospheric CH₄ over the last 800,000 years." *Nature* 453, May 15, 2008, 383–386.
- Lüthi, D., M. Le Floch, B. Bereiter, T. Blunier, J.-M. Barnola, U. Siegenthaler, D. Raynaud, et al. "High-resolution carbon dioxide concentration record 650,000–800,000 years before present." *Nature* 453, May 15, 2008, 379–382.
- Masson-Delmotte, V., A. Landais, N. Combourieu-Nebout, U. Von Grafenstein, J. Jouzel, N. Caillon, J. Chappellaz, D. Dahl-Jensen, S. J. Johnsen, and B. Stenni. "Rapid climate variability during warm and cold periods in polar regions and Europe." *Comptes Rendus Geoscience* 337, 2005, 935–946.
- OCLIB. "Informe de seguimiento del convenio Universitat de les Illes Balears-Conselleria de Medi Ambient del Govern de les Illes Balears, sobre el Observatori del Clima de les Illes Balears." Unpublished technical report. Grup de Meteorologia, UIB, 2007.
- Ruddiman, W. F., ed. *Tectonic Uplift and Climate Change*. New York: Plenum Press, 1997.
- Thompson, D. W., J. J. Kennedy, J. M. Wallace, and P. D. Jones. "A large discontinuity in the mid-twentieth century in observed global-mean surface temperature." *Nature* 453, May 29, 2008, 646–649.
- Wang, Y. M., J. L. Lean, and N. R. Sheeley. "Modeling the sun's magnetic field and irradiance since 1713." *Astrophys. J.* 625, 2005, 522–538.
- Yang, F., and M. Schlesinger. "On the surface and atmospheric temperature changes following the 1991 Pinatubo volcanic eruption: a GCM study." *J. Geophys. Res.-Atmos.* 107, April 2002: doi:10.1029/2001JD000373.

Acknowledgements:

My thanks to Professor C. Ramis, whose suggestions have considerably improved the quality of this text. The figures are exact reproductions of those appearing in IPCC documents and are included here with their tacit authorization.

Global warming: planetary energy balance

The climate depends on exchanges of heat energy between the Sun and the Earth, and between the Earth's surface (and the oceans) and its atmosphere. The planetary (or radiative) energy balance combines the entry and exit of this radiation that produces heat in the Earth's atmosphere. When the climatic system is in equilibrium, these exchanges are balanced (the entry and exit of radiation compensate each other). If the balance is

altered, a heating or cooling of the atmosphere takes place. The hypothesis of global warming by the greenhouse effect signifies that the amount of radiation the Earth returns to outer space is less than that which it receives. When that happens, if conditions are not reversed, the temperature of the Earth's atmosphere begins to rise continuously (as do the mean temperatures of the seas and of the Earth's surface).

Greenhouse effect

The Earth's radiation becomes "trapped" in the lower layers of the atmosphere because of the "greenhouse effect," which is caused by certain gasses—water vapor, carbon dioxide (CO₂), methane, nitrous oxide, and others—that produce an effect similar to that of a greenhouse's glass. That is, they let solar radiation in, but prevent the Earth's heat from getting out. The Earth's atmosphere has always suffered a "natural" greenhouse effect, mainly due to water vapor, and this has kept its mean temperature around 15°C (instead of the -18°C it would otherwise have). That equilibrium is due to the natural evolution of the Earth, and to successive economic revolutions (agricultural, urban, and industrial) undergone by humanity over the course of its history. What was new in the twentieth century, especially in its final decades, was the acceleration of the concentration of greenhouse gasses (GG)—especially CO₂—in the atmosphere. These persistently unbalanced the *planetary energy balance* and marked a growing tendency toward warming.

IPCC

Intergovernmental Panel for Climate Change (IPCC). A group of experts created in 1988 by the World Meteorological Organization (WMO) and the United Nations Environment Programme (UNEP) for the study of climate change. It is organized in three working groups that deal respectively with the scientific aspects of climate and climate change; the impact of change and adaptive measures that must be taken to minimize it; and measures tending to mitigate it. Every six years, the IPCC publishes a report of its work. The latest (4th), published in 2007, has strongly reinforced scientific consensus on global warming by increasing the concentration of greenhouse gasses, and worldwide awareness of the need to take action.

The Kyoto Protocol

In December 1997, the developed countries signed an agreement in Kyoto (Japan) in which they promised to reduce emissions of the main greenhouse gasses by an average of 5.2% over the period 2008–12, with regard to the levels of 1990. Excesses or defects in the reduction of emissions by specific countries could be compensated by an *emissions market* that would organize the purchase and sales of emission rights. The agreement was supposed to enter effect when ratified by countries responsible for at least 55% of CO₂ emissions. Despite reticence by several countries (Japan, Canada, Australia, and, most of all, the United States—the world's leading emitter of GGs), this threshold was crossed in December 2004 following ratification by Russia, and the Protocol went into force in February 2005. Nevertheless, awareness of the need for an agreement involving all developed and emerging countries, along with the worldwide

repercussions of the 4th IPCC Report, Al Gore's documentary, *An Inconvenient Truth*, and the Stern Review, led to a "road map" signed at the last minute by 190 countries at the Bali Conference in December 2007. The signatories are convinced that, in order to avoid the worst effects of global warming, worldwide emissions of greenhouse gasses should reach their maximum in ten to fifteen years and descend drastically from then on. This agreement spreads the effort among developed countries (which must cut their emissions by as much as 40% from 2020) and developing countries, which must follow more sustainable growth strategies. It also foresees financing for measures necessary to adapt to the negative consequences of "compromised" (irreversible) climate change, and for the transfer of "clean" technologies to developing countries. Before the end of 2012, when the Kyoto Protocol ends, a complete agreement must be widely ratified.

The Stern Review

The Government of the United Kingdom commissioned economist, Nicholas Stern, to study the economics of climate change. His findings were published in October 2006 as *The Stern Review on the Economics of Climate Change*. His fundamental thesis is that the cost of strategies to mitigate the effects of climate change may not cost more than 1% of the world's GNP (the maximum would be 3%). On the other hand, if action were not taken immediately, damage caused by global warming would reach an

annual cost of 5% of the worldwide GNP, starting now and lasting forever. In certain circumstances, it could reach 20% of the world's GNP, making it equivalent to the effects of the twentieth century's world wars and the Great Depression. The Stern Review has led to considerable argument about those figures, about the financing of measures of mitigation and adaptation, and about the distribution of the costs of fighting climate change among the present generation and those to come.